IN THE CLAIMS

1. (Original): Compounds having general formula (I)

$$X_1$$
 X_2
 X_3
 X_4
 X_4
 X_4
 X_4
 X_4

wherein:

- a group selected from X₁, X₂ and X₃ represents an R group;
- X₄ and X₅ and two of the remaining X₁, X₂, X_{.3} groups represent a hydrogen atom or a halogen atom, on the condition that at least two of said groups represent a halogen atom;
- R represents a C₁-C₁₂ alkyl or haloalkyl group; a C₁-C₁₂ alkoxy or alkylthio group optionally substituted by halogen atoms, cyano groups, C₁-C₆ alkoxy groups optionally halogenated, C₂-C₁₀ alkoxyalkoxy groups optionally halogenated, NH₂ groups optionally substituted by C₁-C₆ alkyl groups optionally halogenated, C₃-C₁₂ trialkyl silyl groups, aryloxy or heteroaryloxy groups, in turn optionally substituted by halogen atoms, C₁-C₆ alkyl groups optionally halogenated, C₁-C₆ alkoxy groups optionally halogenated, nitro groups, cyano groups; a C₂-C₁₂ alkenyloxy or alkenylthio group optionally substituted by halogen atoms, cyano groups, aryl groups or heteroaryl groups, in turn optionally substituted by halogen atoms, C₁-C₆ alkyl, halo-alkyl, alkoxy, halo-alkoxy groups, nitro groups, cyano groups; a C₃-C₁₂ alkinyloxy or alkinylthio group optionally substituted by halogen atoms, C₁-C₆ alkoxy or haloalkoxy groups, aryl or heteroaryl groups, in turn optionally substituted by halogen atoms, C₁-C₆ alkoxy or haloalkoxy groups, aryl or heteroaryl groups, in turn optionally substituted by halogen atoms, C₁-C₆ alkyl, haloalkyl, alkoxy, halo-

alkoxy groups, nitro groups, cyano groups; a linear or branched C3-C12 alkoxyiminoalkylidenoxy or alkoxyiminoalkylidenthio group; a C3-C8 cycloalkoxy or cycloalkylthio group optionally substituted by halogen atoms, C1-C6 alkyl, haloalkyl, alkoxy, haloalkoxy groups; a C4-C12 cycloalkylalkoxy or cycloalkylalkylthio group optionally substituted by halogen atoms, C1-C6 alkyl, haloalkyl, alkoxy, haloalkoxy groups; a heterocyclyloxy, heterocyclylthio, heterocyclyl-(C1-C6)alkoxy or heterocyclyl-(C1-C6)alkylthio group, optionally substituted by halogen atoms, (C1-C6) alkyl, haloalkyl, alkoxy, haloalkoxy groups; an aryloxy, arylthio, hetero-aryloxy, heteroarylthio, aryl-(C1-C6)alkoxy, aryl-(C1-C6)alkylthio, heteroaryl-(C1-C6)alkoxy or heteroaryl-(C1-C6)alkylthio group optionally substituted by halogen atoms, C1-C6 alkyl groups optionally halogenated, C1-C6 alkoxy groups optionally halogenated, nitro groups, cyano groups;

- A, the same or different, when n is greater than or equal to 2, represents a halogen atom or a C₁-C₄ alkyl, haloalkyl, alkoxyl, haloalkoxyl group;
- Y represents an OCH3 group, an NHCH3 group, an NH2 group;
- Z represents a CH group or a nitrogen atom N;
- n is an integer ranging from 0 to 4.
 - 2. (Original): The compounds according to claim 1, characterized in that they are an isomeric mixture in any proportion, or the isomer E or the isomer Z of the compounds having formula (f).
 - 3. (Original): The compounds according to claim 1, characterized in that they are the isomer E of the compounds having formula (I).
 - 4. (Original): The compounds according to claim 1, characterized in that X₃ represents an R group according to the above mentioned meanings, X2 and X4 represent a halogen atom, X1 and X5 represent a hydrogen atom and n is equal to 0.

- 5. (Original): The compounds according to claim 1, characterized in that they are selected from:
- methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxymethyl)phenyl]-3-methoxyacrylate;
- methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxymethyl)phenyl]-2-methoxyiminoacetate;
- (E)-2-[2-(4-cyclopropylmethoxy-3,5-dichlorophenoxymethyl)phenyl]-N-methyl- 2-methoxyiminoacetamide;
- methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxymethyl]ph-enyl}-3-methoxyacrylate;
- methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxymethyl]phenyl}-2-methoxyiminoacetate;
- (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxymethyl]phenyl}-N-methyl-2-methoxyiminoacetamide;
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxymethyl]phenyl-}-3-methoxyacrylate;
- methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxymethyl]phenyl-}-2-methoxyiminoacetate;
- (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)-phenoxymethyl]pheny-1}-N-methyl-2-methoxyminoacetamide;
- methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxyme-thyl]phenyl}-3-methoxyacrylate; methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxymethyl]phenyl}-2-methoxyiminoacetate;
- (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxymethyl]phenyl}-N-methyl-2-methoxyiminoacetamide;
- methyl (E)-2-[2-(4-cyclobutylmethoxy-3,5-dichlorophenoxymethyl)phenyl]-3-methoxy- acrylate; methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dimethylbutoxy)phenoxymethyl]phenyl}-3-meth- oxyacrylate; methyl (E)-2-{2-[3,5-dichloro-4-(3-methylbutoxy)phenoxymethyl]phenyl}-3-methoxyacrylate; methyl (E)-2-[2-(4-cyclohexylmethoxy-3,5-dichlorophenoxymethyl]phenyl}-3-

- methyl (E)-2-{2-[3,5-dichloro-4-(2,4-ichlorobenzyloxy)phenoxymethyl]phenyl}-3-m- ethoxyacrylate; methyl (E)-2-{2-[3,5-dichloro-4-(4-chlorobenzyloxy)phenoxymethyl]phenyl}-3-metho-xyacrylate.
- 6. (Original): The process for the preparation of the compounds having general formula (I), according to any of the claims 1-5, characterized in that it includes a condensation reaction of a compound having general formula (II) with a phenol having general formula (III), according to the reaction scheme 1: wherein, X_1 , X_2 , X_3 , X_4 and X_5 , A, Y, Z and n have the meanings defined above, L represents a leaving group such as a chlorine atom, a bromine atom or a R_1SO_3 group wherein R_1 represents a C_1 - C_6 alkyl or haloalkyl, or a phenyl optionally substituted.
- 7. (Original): The process according to claim 6, characterized in that the reaction is carried out in an inert organic solvent, at a temperature ranging from 0° C and the boiling temperature of the reaction mixture, possibly in the presence of an inorganic or organic base.
- 8. (Original): The process according to claim 7, characterized in that the solvent is selected from alcohols, ethers, esters, ketones, chlorinated hydrocarbons, aromatic hydrocarbons, aliphatic hydrocarbons, aprotic dipolar solvents.
- 9. (Original): The process according to claim 7, characterized in that the inorganic base is selected from hydrides, hydroxides, carbonates of alkaline or alkalineerath metals.
- 10. (Original): The process according to claim 7, characterized in that the organic base is selected from pyridine, dimethylaminopyridine, aliphatic amines, cyclic amines, alcoholates of alkaline metals.
- 11. (Original): Use of the compounds having general formula (I) wherein: a group

selected from X1, X2 and X3 represents an R group; X4 and X5 and two of the remaining X1, X2, X3 groups represent a hydrogen atom or a halogen atom, on the condition that at least two of said groups represent a halogen atom; R represents a C1-C12 alkyl or haloalkyl group; a C1-C12 alkoxy or alkylthio group optionally substituted by halogen atoms, cyano groups, C1-C6 alkoxy groups optionally halogenated, C2-C10 alkoxyalkoxy groups optionally halogenated, NH2 groups optionally substituted by C1-C6 alkyl groups optionally halogenated, C3-C12 trialkyl silyl groups, aryloxy or heteroaryloxy groups, in turn optionally substituted by halogen atoms, C1-C6 alkyl groups optionally halogenated, C1-C6 alkoxy groups optionally halogenated, nitro groups, cyano groups; a C2-C12 alkenyloxy or alkenylthio group optionally substituted by halogen atoms, cyano groups, aryl groups or heteroaryl groups, in turn optionally substituted by halogen atoms, C1-C6 alkyl, haloalkyl, alkoxy, halo-alkoxy groups, nitro groups, cyano groups; a C3-C12 alkinyloxy or alkinylthio group optionally substituted by halogen atoms, C1-C6 alkoxy or haloalkoxy groups, aryl or heteroaryl groups, in turn optionally substituted by halogen atoms, C1-C6 alkyl, haloalkyl, alkoxy, haloalkoxy groups, nitro groups, cyano groups; a linear or branched C3-C12 alkoxyiminoalkylidenoxy or alkoxyiminoalkylidenthio group; a C3-C8 cycloalkoxy or cycloalkylthio group optionally substituted by halogen atoms, C1-C6 alkyl, haloalkyl, alkoxy, halo-alkoxy groups; a C4-C12 cycloalkylalkoxy or cycloalkylalkylthio group optionally substituted by halogen atoms, C1-C6 alkyl, haloalkyl, alkoxy, halo-alkoxy groups; a heterocyclyloxy, heterocyclylthio, heterocyclyl(C₁-C₆)alkoxy or hetero-cyclyl(C₁-C₆)alkylthio group, optionally substituted by halogen atoms, (C1-C6) alkyl, haloalkyl, alkoxy, halo-alkoxy groups; an aryloxy, arylthio, beteroaryloxy, heteroarylthio, aryl-(C1-C6)alkoxy, aryl-(C1-C6)alkylthio, heteroaryl-(C1-C6)alkoxy or heteroaryl-(C1-C6)alkylthio group optionally substituted by halogen atoms, C₁-C₆ alkyl groups optionally halogenated, C1-C6 alkoxy groups optionally halogenated, nitro groups, cyano groups; A, the same or different, when n is greater than or equal to 2, represents a halogen atom or a C₁-C₄ alkyl, haloalkyl, alkoxyl, haloalkoxy group; Y represents an OCH₃ group, an NHCH₃ group, an NH₂ group; Z represents a CH group or a

nitrogen atom N; n is an integer ranging from 0 to 4; as acaricides and/or insecticides and/or fungicides.

- 12. (Original): The use according to claim 11 of the isomers E of the compounds having formula (I).
- 13. (Original): The use according to claim 11, wherein X.sub.3 represents an R group according to the above meanings, X_2 and X_4 represent a halogen atom, X_1 and X_5 represent a hydrogen atom and n is equal to 0.
- 14. (Original): The use according to claim 11, wherein the compounds of formula (I) are selected from: methyl (E)-2-[2-(4-cyclopropylmethoxy-3,5dichlorophenoxymcthyl)phenyl]-3-methox- yacrylate; methyl (E)-2-[2-(4cyclopropylmethoxy-3,5-dichlorophenoxymethyl)phenyl]-2-methoxyiminoacetate; (E)-2-[2-(4-cyclopropylmcthoxy-3,5dichlorophenoxymethyl)phenyl]-N-methyl--2-methoxyiminoacetamide; methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxymethyl]phenyl}-3-methoxyacrylate; methyl (E)-2-{2-[4-(2,2-dichlorocyclopropyl)methoxy-3,5-dichlorophenoxymethyl]ph-enyl}-2-methoxyiminoacetate; (E)-2-{2-[4-(2,2dichlorocyclopropyl)methoxy-3,5-dichlorphenoxymethyl]phenyl}-N-methyl-2methoxyiminoacetamide; methyl (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2enyloxy)phenoxymethyl]phenyl-}-3-methoxyacrylate; methyl (E)-2-{2-[3,5dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxymethyl]phenyl- }-2methoxyiminoacetate; (E)-2-{2-[3,5-dichloro-4-(3,3-dichloroprop-2-enyloxy)phenoxymethyl]pheny-1}-N-methyl-2-methoxyiminoacetamide; methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxyme-thyl]phenyl}-3-methoxyacrylate; methyl (E)-2-{2-[3,5-dichloro-4-(3-chloro-4,4,4-trifluorobut-2-enyloxy)phenoxyme-thyl]phenyl}-2-methoxyiminoacetate; (E)-2-{2-[3,5dichloro-4-(3-chloro-4,4,4-tri-fluorobut-2-enyloxy)phenoxym- ethyl]phenyl}-Nmethyl-2-methoxyiminoacetamide; methyl (E)-2-[2-(4-cyclobutylmethoxy-3,5dichlorophenoxymethyl)phenyl]-3-methoxy- acrylate; methyl (E)-2-{2-[3,5-

dichloro-4-(3,3-dimethylbutoxy)phenoxymethyl]phenyl}-3-meth- oxyacrylate; methyl (E)-2-{2-[3,5-dichloro-4-(3-methylbutoxy)phenoxymethyl]phenyl}-3-methoxya- crylate; methyl (E)-2-[2-(4-cyclohexylmethoxy-3,5-dichlorophenoxymethyl]phenyl}-3-methoxy- acrylate; methyl (E)-2-{2-[3,5-dichloro-4-(2,4-dichlorobenzyloxy)phenoxymethyl]phenyl}-3-m- ethoxyacrylate; methyl (E)-2-{2-[3,5-dichloro-4-(4-chlorobenzyloxy)phenoxymethyl]phenyl}-3-metho- xyacrylate.

- 15. (Original): The use according to any of the claims 11-14 for the control of adults, larvae and eggs of mites and insects which are harmful in the agrarian, civil and zoo-technical field.
- 16. (Original): The use according to claim 15, wherein the harmful mites and/or insects are tetranychidae (Tetranychus urticae, Tetranychus telarius, Tetranychus cinnabarinus, Eotetranychus carpini, Panonychus ulmi, Panonychus citri, etc.), eriophyidae (Phytoptus avellanae, Eriophyes vitis, Eriophyes piri, etc.) tarsonemidae (Steneotarsonemus pallidus, etc.), hemiptera (Macrosiphum cuphorbiae, Aphis fabac, Myzus persicae, etc.), lepidoptera (Spodoptera spp., Heliothis spp., Chilo spp., Carpocapsa pomonella, etc.), coleoptera (Leptinotarsa decemlineata, Phaedon cochleariae, etc.), diptera (Aedes spp., Culex spp., Musca spp., etc.).
- 17. (Original): The use according to any of the claims 11-14 for the control of phytopathogenous fungi such as: Helminthosporium spp., Erysiphe spp., Puccinia spp., Plasmopara viticola, Pythium spp., Phytophthora spp., Rhynchosporium spp., Septoria spp., Sphaerotheca fuliginea, Podosphaera leucotricha, Pyricularia oryzae, Uncinula necator, Venturia spp., Botrytis cinerea, Fusarium spp., Alternaria spp., Cercospora spp.
- 18. (Original): The use according to any of the claims 11-14 for the control of mites, insects and fungi which are harmful in crops of agrarian and horticultural

interest, on domestic and breeding animals, in environments frequented by human beings.

- 19. (Original): A method for controlling mites and/or insects and/or phytopathogenous fungi in crops of agrarian and horticultural interest, and/or on domestic and breeding animals, and/or in environments frequented by human beings, by the application of the compounds having general formula (I) according to one of the claims 1-5.
- 20. (Original): The method according to claim 19, characterized in that the quantity of compound to be applied varies from 10 g to 5 kg per hectare.
- 21. (Original): Acaricidal and/or insecticidal and/or fungicidal compositions containing as active principle one or more compounds having general formula (I) according to one of the claims 1-5.
- 22. (Original): The compositions according to claim 21, comprising other active principles compatible with the compounds having general formula (I), such as other acaricides/insecticides, fungicides, phyto-regulators, antibiotics, herbicides, fertilizers.
- 23. (Original): The compositions according to claim 21, characterized in that the concentration of active principle ranges from 1 to 90%, preferably from 5 to 50%.